



EU 1737-132

Brandt, Søren; Castro-Tirado, A. J.; Lund, Niels

Published in:
International Astronomical Union Circulars (IAUC)

Publication date:
1992

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Brandt, S., Castro-Tirado, A. J., & Lund, N. (1992). EU 1737-132. *International Astronomical Union Circulars (IAUC)*, (5643). <http://www.cbat.eps.harvard.edu/iauc/05600/05643.html#Item1>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Central Bureau for Astronomical Telegrams
 INTERNATIONAL ASTRONOMICAL UNION
 Postal Address: Central Bureau for Astronomical Telegrams
 Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
 Telephone 617-495-7244/7440/7444 (for emergency use only)
 TWX 710-320-6842 ASTROGRAM CAM EASYLINK 62794505
 MARSDEN@CFA or GREEN@CFA (.SPAN, .BITNET or .HARVARD.EDU)

EU 1737-132

S. Brandt, A. J. Castro-Tirado, and N. Lund report: "A new x-ray transient in Serpens Cauda has been discovered by the WATCH wide-field monitor on the ESA EURECA satellite. The transient was first observed on Oct. 11, at R.A. = 17h37m20s, Decl. = -13 15'.0 (equinox 1950.0; error radius 45'). The source brightness at the time of discovery was about 500 mCrab (above 6 keV), and remained at about this level at least until Oct. 18. In the spectral band between 6 and 15 keV, the source is now only slightly harder than that of Sco X-1, though during the first two days of the outburst the spectrum was harder. Our limited sensitivity does not allow us to detect the source above 15 keV. The source may be similar to that of dwarf x-ray transients such as N Mus 1991 and GRO J0422+32. Follow up at other wavelengths is encouraged."

PERIODIC COMET SWIFT-TUTTLE (1992t)

P. Colom, D. Bockelee-Morvan, G. Bourgois, J. Crovisier, E. Gerard, and L. Jorda, Paris-Meudon Observatory, communicate: "The main lines of the OH radical at 1667 and 1665 MHz were detected in emission in P/Swift-Tuttle on Oct. 15 with the Nancay Radio Telescope. For the period Oct. 16-20, the 1667-MHz profile has a peak flux density of 86 +/- 14 mJy, a FWHM of 2 km/s, and is blueshifted by 0.3 km/s with respect to the radial velocity of the nucleus. The line area of 184 +/- 17 mJy km/s corresponds to an OH production rate of 8×10^{28} molecules/s."

J. V. Scotti, University of Arizona, reports that CCD images were obtained of this comet on Oct. 18.1, 19.1, and 21.1 UT with the Spacewatch telescope, showing an ion tail extending at least 15' toward the north-northeast. There was a noticeable increase in inner coma activity, as well as in overall brightness, between Oct. 18.1 and 19.1. On Oct. 21.097 UT, there was a 10'.5 coma surrounding an inner coma that is asymmetrical toward the west (though the overall coma is nearly circular). A fan-shaped jet to the west dissipated slightly between Oct. 18 and 21, and the tail was a bit fainter on Oct. 21 than on previous nights. The fan-shaped jet structure on Oct. 21 runs from p.a. 242 to 313 deg; the main tail extends toward p.a. 24 deg, and there are two faint streamers bracketing the tail at p.a. 9 and 50 deg, each extending about 5' and perhaps further.